

# Emergyl

## Description:

Giloy, Amalaki, Haritaki are called the RASAYANA in the ayurveda by virtue of its antioxidant action scavenges free radicals and its immunomodulatory activity enhances body immunity. Emergyl Powder helps delay the physiological changes associated with aging, revives physical capacity, raises the threshold of fatigue and promotes well-being. Along with Shatavari, Ashwagandha, Vidarikand, Agastya, Akkalkara, Umbar, Rewandachini it helps to rejuvenate the body tissues and is the natural source of energy. It accelerates cellular regeneration and repair. Marich, Saalam, Tulsi, Kantakari, Neem, Punnag of Emergyl Powder facilitates respiratory functions. Emergyl Powder regulates fat and carbohydrate metabolism by the carminative, digestive, stimulant actions of Chitrak, Vidang, Shigru, Marich, Bhumi amla and Gokshur, and improves appetite, digestion and assimilation. Dadim, Arjuna, Bharangi are looking after the health of the heart, assists cardiovascular functioning by improving circulation and reducing raised lipid levels. So Emergyl Powder rejuvenates the entire body system with Kewanch beej, Lajjawanti, Yashtimadhu, Jaifal and retards the skin changes associated with aging.

## Composition:

Each 500 g contains :

Amruta (*Tinospora Cordifolia*) - 260 gm.

*Sisbania grandiflora*, *Anacyclus pyrethrum*, *Ficus glomerata*, *Piper nigrum*, *Rheum emodi*, *Mimosa pudica*, *Colophyllum inophyllum*, *Moringa pterygosperma*, *Myristica fragrans*, *Embelia ribes*, *Clerodendron serratum*, *Plumbago zeylanica*, *Andrographis paniculata*, *Terminalia arjuna*, *Punica granatum*, *Tribulus terrestris*, *Pueraria tuberosa* 5 gm each.  
*Solanum xanthocarpum*, *Mucuna pruriens*, *Boerhavia diffusa*, *Orchis mascula* o *latifolia*, *Ocimum sanctum*, *Terminalia chebula*, *Embllica officinalis*, *Asparagus racemosus* - 10 gm. each  
*Azadirachta indica*, *Phyllanthus niruri*, *Withania somnifera* -15 gm. each  
*Glycyrrhiza glabra* -12.5 gm.

## Indication:

Balya tonic. Regenerative for patient with health conditions : Severe infection, Chronic fever, Typhoid, Malaria, Tuberculosis with/without lymphadenopathy, AIDS. Convalescence, Fatigue...

## Dosage:

- In chronic cases : 5 gm, 2 times a day along with water/milk after meals.
  - In preventive cases : 5 gm, once a day along with water/milk after meals.
  - In life threatening conditions : 5 gm, 3 times a day along with water/milk after meals.
- Or as directed by the physician.

## Direction for use:

Mix the required quantity as per dose of powder with water or milk. Prepare in round pill form for easy swallowing. Leave in open for 5 minutes and then swallow it whole.

## Note:

Body weight to be taken before using the medicine and should be monitored carefully, increase in body weight is the 1st indication of positive effect of the medicine. Do not smoke, consume liquor/tobacco during therapy.

## Storage:

Keep the lid tightly closed out of reach of children. Keep in a cool dry place away from direct sunlight. Do not use if seal is broken. Use spoon provided.